

FLOER HOMOLOGY OF BRIESKORN HOMOLOGY SPHERES : SOLUTION TO ATIYAH'S PROBLEM

NIKOLAI SAVELIEV

ABSTRACT. In this paper we answer the question posed by M. Atiyah, see [12], and give an explicit formula for Floer homology of Brieskorn homology spheres in terms of their branching sets over the 3-sphere. We further show how Floer homology is related to other invariants of knots and 3-manifolds, among which are the $\bar{\mu}$ -invariant of W. Neumann and L. Siebenmann and the Jones polynomial. Essential progress is made in proving the homology cobordism invariance of our own ν -invariant, see [15].

Let p, q , and r be pairwise coprime positive integers. The Brieskorn homology 3-sphere $\Sigma(p, q, r)$ is the link of the singularity of $f^{-1}(0)$, where $f : \mathbb{C}^3 \rightarrow \mathbb{C}$ is a map of the form $f(x, y, z) = x^p + y^q + z^r$. The complex conjugation in $\mathbb{C}^3$ acts on $\Sigma(p, q, r)$ turning it into a double branched cover of S^3 branched over a Montesinos knot $k(p, q, r)$. In this paper, we express the Floer instanton homology groups $I_n(\Sigma(p, q, r))$, see [5], in terms of the Casson's λ -invariant of $\Sigma(p, q, r)$ and the signature of the knot $k(p, q, r)$.

Theorem 1. *The Floer homology group $I_n(\Sigma(p, q, r))$ is trivial if n is odd, and is a free abelian group of the rank*

$$\text{rank } I_n(\Sigma(p, q, r)) = \frac{1}{16} (8\lambda(\Sigma(p, q, r)) - (-1)^{n/2} \text{sign } k(p, q, r))$$

if n is even.

The groups $I_*(\Sigma(p, q, r))$ were studied by R. Fintushel and R. Stern in [6] (we use the Floer index convention of this paper). They gave an algorithm which for any given set of positive coprime integers p, q, r computes the ranks of $I_*(\Sigma(p, q, r))$. However, their approach does not give a closed form formula for $I_*(\Sigma(p, q, r))$. It should be mentioned though that the vanishing of the groups $I_n(\Sigma(p, q, r))$ for odd n follows from [6]; we give an alternative proof in Section 3.

1991 *Mathematics Subject Classification.* 57M25, 57R80.

Key words and phrases. Floer homology, Seifert manifolds, Montesinos knots, Casson invariant, Jones polynomial, homology cobordism, knot signature.

Research at MSRI is supported in part by NSF grant DMS-9022140.

In [15] we introduced an invariant ν , which for Brieskorn homology spheres takes the form

$$\nu(\Sigma(p, q, r)) = \frac{1}{2} \sum_{n=0}^3 (-1)^{n+1} \text{rank } I_{2n}(\Sigma(p, q, r)). \quad (1)$$

Note that if the ranks in (1) were added without plus or minus signs one would get the Casson invariant $\lambda(\Sigma(p, q, r))$. We conjectured in [15] that the ν -invariant equals the $\bar{\mu}$ -invariant of W. Neumann [13] and L. Siebenmann [17] for all Seifert fibered homology spheres. We prove that our conjecture is true. The first step is given by the following theorem.

Theorem 2. *For any Brieskorn homology 3-sphere $\Sigma(p, q, r)$, the ν -invariant defined by (1) and the $\bar{\mu}$ -invariant coincide, $\nu(\Sigma(p, q, r)) = \bar{\mu}(\Sigma(p, q, r))$.*

It can be easily seen that Theorem 2 is in fact equivalent to Theorem 1 due to the following two observations: first, $\bar{\mu}(\Sigma(p, q, r)) = 1/8 \cdot \text{sign } k(p, q, r)$, see [17], and second, $I_n(\Sigma(p, q, r)) = I_{n+4}(\Sigma(p, q, r))$ for all n . Second step in proving our conjecture makes use of the so called splicing additivity proven for both the $\bar{\mu}$ - and ν -invariants in [15]. Namely, $\bar{\mu}(\Sigma(a_1, \dots, a_n)) = \bar{\mu}(\Sigma(a_1, \dots, a_j, p)) + \bar{\mu}(\Sigma(q, a_{j+1}, \dots, a_n))$, where j is any integer between 2 and $n-2$, and the integers $q = a_1 \cdots a_j$ and $p = a_{j+1} \cdots a_n$ are the products of the first j and the last $(n-j)$ Seifert invariants, respectively. The same additivity holds for the invariant ν .

Theorem 3. *For any Seifert fibered homology 3-sphere $\Sigma(a_1, \dots, a_n)$, the invariants ν and $\bar{\mu}$ coincide, $\nu(\Sigma(a_1, \dots, a_n)) = \bar{\mu}(\Sigma(a_1, \dots, a_n))$.*

Another reformulation of Theorem 1 establishes links between Floer homology and Jones polynomial. The result below follows from Theorem 1 and D. Mullins formula for the Casson invariant of two-fold branched covers [11].

Theorem 4. *Let $\Sigma(p, q, r)$ be a Brieskorn homology sphere then*

$$\text{rank } I_0(\Sigma(p, q, r)) = -\frac{1}{12} \cdot \left. \frac{d}{dt} \right|_{t=-1} \ln V_k(t)$$

where V_k is the Jones polynomial of the Montesinos knot $k = k(p, q, r)$.

Theorem 3 has the following application to the homology 4-cobordism. In [15] and [16] we proved that $\bar{\mu}$ is a homology cobordism invariant for some classes of Seifert fibered homology 3-spheres. The corresponding results for the ν -invariant hold due to the identification $\bar{\mu} = \nu$ in Theorem 3.

Theorem 5. (1) *Let $\Sigma(a_1, \dots, a_n)$ be a Seifert fibered homology sphere homology cobordant to zero. Then $\nu(\Sigma(a_1, \dots, a_n)) \geq 0$.*

(2) *Let $\Sigma = \Sigma(p, q, pqm \pm 1)$ be a surgery on a (p, q) -torus knot, and suppose that Σ is homology cobordant to zero. Then $\nu(\Sigma) = 0$.*

(3) *For all known Seifert fibered homology 3-spheres Σ which are known to be homology cobordant to zero including the lists of Casson-Harer [3] and Stern [18], $\nu(\Sigma) = 0$.*

We will prove that $\nu(\Sigma(p, q, r)) = 1/8 \cdot \text{sign } k(p, q, r)$, which is equivalent to proving Theorem 1. The idea is shortly as follows. In [19], C. Taubes proved that, for any homology 3-sphere Σ ,

$$\lambda(\Sigma) = 1/2 \cdot \chi(I_*(\Sigma)). \quad (2)$$

The Casson's λ -invariant on the left is defined topologically using a Heegaard splitting of Σ and $\text{SU}(2)$ -representation spaces. The number on the right is the Euler characteristic of Floer homology $I_*(\Sigma)$; it can be defined using $\text{SU}(2)$ gauge theory as an infinite dimensional generalization of the classical Euler characteristic.

Let now $\Sigma = \Sigma(p, q, r)$ be endowed with the involution σ induced on $\Sigma \subset \mathbb{C}^3$ by the complex conjugation. We work out σ -invariant versions of both invariants in (2) in this particular situation. We use a σ -invariant Heegaard splitting of Σ and the corresponding σ -invariant $\text{SU}(2)$ -representation spaces to define an invariant $\lambda^\rho(\Sigma)$ in a manner similar to that of A. Casson. On the other hand, a σ -invariant gauge theory produces a σ -invariant Euler characteristic which we denote $\chi^\rho(\Sigma)$. Note that the latter can be defined without actually working out any Floer homology. Then, a σ -invariant version of the Taubes result (2) is that

$$\lambda^\rho(\Sigma(p, q, r)) = 1/2 \cdot \chi^\rho(\Sigma(p, q, r)). \quad (3)$$

Our next step is to show that $\lambda^\rho(\Sigma(p, q, r)) = 1/8 \cdot \text{sign } k(p, q, r)$. We achieve this by pushing σ -invariant representations down to the quotients. We do this for all the manifolds in a σ -invariant Heegaard splitting and note that the push-down representations are only defined on the complements of the fixed point sets, and they map all the meridians to trace-free matrices in $\text{SU}(2)$. After that we are in position to identify our λ^ρ -invariant with the Casson-Lin invariant $1/8 \cdot \text{sign } k(p, q, r)$ of [8]. The crucial observation is that *all* the representations of $\pi_1(\Sigma(p, q, r))$ are in fact σ -invariant, and the λ^ρ -invariant just counts them with signs different from those defined by Casson for his λ -invariant.

Finally, we identify $1/2 \cdot \chi^\rho(\Sigma)$ with $\nu(\Sigma)$ as follows. We choose a σ -invariant metric on Σ and an almost complex structure on $\Sigma \times \mathbb{R}$ so that the involution $\sigma \times 1$ on $\Sigma \times \mathbb{R}$ is anti-holomorphic. At the level of tangent spaces, the ± 1 -eigenspaces of $\sigma \times 1$ are then isomorphic to each other and hence have the same dimension. Therefore, the spectral flow used in [19] to define $\chi(\Sigma)$ splits in halves, and the ν -invariant becomes an “honest” σ -invariant Euler characteristic equal to $1/2 \cdot \chi^\rho(\Sigma)$.

The paper begins with an introduction to the relevant topology of Brieskorn homology spheres and Montesinos knots. In Section 2 the invariant $\lambda^\rho(\Sigma(p, q, r))$ is introduced and the equality $\lambda^\rho(\Sigma(p, q, r)) = 1/8 \cdot \text{sign } k(p, q, r)$ is proven. The invariant

$\chi^\rho(\Sigma(p, q, r))$ is defined in Section 3; the equality $1/2 \cdot \chi^\rho(\Sigma(p, q, r)) = \nu(\Sigma(p, q, r))$ is proven in the same section. Section 4 is devoted to the proof of the fact that $\lambda^\rho(\Sigma(p, q, r)) = 1/2 \cdot \chi^\rho(\Sigma(p, q, r))$. Each section has its own introduction and is further subdivided.

It should be pointed out that most of the results in the paper hold in a more general situation. A description of the relevant results will appear elsewhere, as will a more detailed version of this preprint.

1. TOPOLOGY OF BRIESKORN HOMOLOGY SPHERES

Let p, q, r be relatively prime positive integers greater than or equal to 2. The Brieskorn homology sphere $\Sigma(p, q, r)$ is defined as the algebraic link

$$\Sigma(p, q, r) = \{ (x, y, z) \in \mathbb{C}^3 \mid x^p + y^q + z^r = 0 \} \cap S_1^5,$$

where S_1^5 is the unit sphere in $\mathbb{C}^3$. This is a smooth naturally oriented 3-manifold with $H_*(\Sigma(p, q, r)) = H_*(S^3)$. Moreover, $\Sigma(p, q, r)$ is Seifert fibered, see [14], with the Seifert invariants $\{b, (p, b_1), (q, b_2), (r, b_3)\}$ such that

$$b_1qr + b_2pr + b_3pq + bpqr = 1. \quad (4)$$

The complex conjugation on $\mathbb{C}^3$ obviously acts on $\Sigma(p, q, r)$ as

$$\sigma : \Sigma(p, q, r) \rightarrow \Sigma(p, q, r), \quad (x, y, z) \mapsto (\bar{x}, \bar{y}, \bar{z}). \quad (5)$$

The fixed point set of this action is never empty. The quotient of $\Sigma(p, q, r)$ by the involution σ is S^3 , with the branching set the so called Montesinos knot $k(p, q, r)$, see [10], [2], or [17]. The knot $k(p, q, r)$ can be described by the following diagram




Figure 1

where a box with α, β in it stands for a rational (α, β) -tangle, see [2], Fig. 12.9. The parameters $b, (p, b_1), (q, b_2), (r, b_3)$ are the Seifert invariants of the corresponding $\Sigma(p, q, r)$. According to [2], Theorem 12.28, these parameters together with (4) determine the knot $k(p, q, r)$ uniquely up to isotopy.

2. THE INVARIANT λ^ρ

In this section we first shortly recall the definition of Casson's λ -invariant. Our λ^ρ -invariant for $\Sigma(p, q, r)$ will be a σ -invariant version of λ . To define it, we introduce a σ -invariant Heegaard splitting of $\Sigma(p, q, r)$. Then we define the corresponding σ -invariant representation spaces and investigate closely the representation space of $\pi_1(\Sigma(p, q, r))$. It turns out that all the representations in the latter space are σ -invariant. After computing the necessary dimensions and checking the transversality condition, we define the λ^ρ -invariant as an intersection number of σ -invariant representation spaces. Finally, we check that $\lambda^\rho(\Sigma(p, q, r)) = 1/8 \cdot \text{sign } k(p, q, r)$.

1. Casson invariant. Let M be an oriented homology 3-sphere with a Heegaard splitting $M = M_1 \cup M_2$ where M_1 and M_2 are handlebodies of genus $g \geq 2$ glued along their common boundary, a Riemann surface M_0 . Let

$$\mathcal{R}(M_i) = \text{Hom}^*(\pi_1(M_i), \text{SU}(2)) / \text{ad } \text{SU}(2), \quad i = 0, 1, 2, \emptyset,$$

be the set of conjugacy classes of irreducible representations of $\pi_1(M_i)$ is $\text{SU}(2)$. Each $\mathcal{R}(M_i)$, $i = 0, 1, 2$, is naturally an oriented manifold. The dimension of $\mathcal{R}(M_1)$ and $\mathcal{R}(M_2)$ is $3g-3$, and that of $\mathcal{R}(M_0)$ is $6g-6$. The inclusions $M_0 \subset M_i$, $i = 1, 2$, induce embeddings $\mathcal{R}(M_i) \subset \mathcal{R}(M_0)$. The points of intersection of $\mathcal{R}(M_1)$ with $\mathcal{R}(M_2)$ are in one-to-one correspondence with $\mathcal{R}(M)$. If the intersection is transversal we define the Casson's λ -invariant as

$$\lambda(M) = \frac{1}{2} \sum_{\alpha \in \mathcal{R}(M)} \varepsilon_\alpha$$

where $\varepsilon_\alpha = \pm 1$ is a sign obtained by comparing the orientations on $T_\alpha \mathcal{R}(M_1) \oplus T_\alpha \mathcal{R}(M_2)$ and $T_\alpha \mathcal{R}(M_0)$. Note that the intersection is transversal if $\Sigma = \Sigma(p, q, r)$.

2. The invariant Heegaard splitting. We are going to construct an invariant Heegaard splitting of $\Sigma(p, q, r)$. Let us first fix notations. By Σ we denote a Brieskorn homology sphere $\Sigma(p, q, r)$ and by $\sigma : \Sigma \rightarrow \Sigma$ the involution constructed in (5). The projection on the quotient space will be denoted by π , so $\pi : \Sigma \rightarrow \Sigma/\sigma = S^3$. The projection π maps the fixed point set $\text{Fix}(\sigma) \subset \Sigma$ onto the Montesinos knot $k = k(p, q, r) \subset S^3$.

The knot $k \subset S^3$ can be represented as the closure of a braid β on n strings. We think about k as consisting of the braid β and n untangled arcs in S^3 forming its closure, see Figure 2. Let $S \subset S^3$ be an embedded 2-sphere in S^3 splitting S^3 in

two 3-balls, B_1 and B_2 , with common boundary S , and such that the entire braid β belongs to $\text{int } B_1$. The intersection $k \cap S$ consists of $2n$ points $P_1, \dots, P_{2n}$.




Figure 2

Now, we define a Heegaard splitting $\Sigma = M_1 \cup_{M_0} M_2$ as follows:

$$M_1 = \pi^{-1}(B_1), \quad M_2 = \pi^{-1}(B_2), \quad M_0 = \pi^{-1}(S). \quad (6)$$

Obviously, M_1 and M_2 are handlebodies branched over the braid β and the n untangled arcs, respectively. Their common boundary is M_0 , which is a closed surface branched over the points $P_1, \dots, P_{2n} \in S$. The genus g of M_0 can be figured out by comparing Euler characteristics,

$$2 \cdot \chi(S) = \chi(M_0) + \chi(\text{Fix}(\sigma) \cap S),$$

where $\chi(S) = 2$, $\chi(\text{Fix}(\sigma) \cap S) = 2n$, and $\chi(M_0) = 2 - 2g$. Therefore, $g = n - 1$. The Heegaard splitting we just defined is σ -invariant in the sense that $\sigma(M_i) = M_i$ for $i = 0, 1, 2$.

3. Representations of Brieskorn homology spheres. Let $\Sigma(p, q, r)$ be a Brieskorn homology 3-sphere, and

$$\mathcal{R}(\Sigma(p, q, r)) = \text{Hom}^*(\pi_1(\Sigma(p, q, r)), \text{SU}(2)) / \text{ad SU}(2) \quad (7)$$

the space of the conjugacy classes of irreducible representations of its fundamental group in $\text{SU}(2)$. In this subsection we are concerned with describing the space $\mathcal{R}(\Sigma(p, q, r))$ and the involution σ^* induced on the representation space by σ .

We will follow [6] and first identify the group $SU(2)$ with the group S^3 of unit quaternions in the usual way, so that

$$i = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad j = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad k = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}.$$

Under this identification, the trace $\text{tr}(A)$ of an element $A \in SU(2)$ coincides with the number $2 \text{Re } A$, $A \in S^3$, and the mapping $r : S^3 \rightarrow [0, \pi]$, $r(A) = \arccos(\text{Re } A)$, is a complete invariant of the conjugacy class of the element A . This conjugacy class is a copy of S^2 in S^3 unless $A = \pm 1$, in which case the conjugacy class consists of just one point.

The fundamental group $\pi_1(\Sigma(p, q, r))$ has the following presentation, see [2],

$$\pi_1(\Sigma(p, q, r)) = \langle x, y, z, h \mid h \text{ central}, x^p = h^{-b_1}, y^q = h^{-b_2}, z^r = h^{-b_3}, xyz = h^{-b} \rangle.$$

Specifying an irreducible representation $\alpha : \pi_1(\Sigma(p, q, r)) \rightarrow SU(2)$ amounts to specifying a set $\{\alpha(h), \alpha(x), \alpha(y), \alpha(z)\}$ of unit quaternions. In fact, we only need to specify the first three quaternions in this set because $\alpha(z)$ will then be expressed in their terms as $\alpha(z) = (\alpha(x)\alpha(y))^{-1}\alpha(h)^{-b}$. Since h is central and the representation α is irreducible, $\alpha(h) = \pm 1$. Let us denote $\varepsilon_i = \alpha(h)^{b_i} = \pm 1$, $i = 1, 2$, and $\varepsilon_3 = \alpha(h)^{b_3 - rb} = \pm 1$. Then the relations $x^p = h^{-b_1}$, $y^q = h^{-b_2}$ and $(xy)^r = h^{b_3 - rb}$ imply the following restrictions on $\alpha(x)$ and $\alpha(y)$:

$$r(\alpha(x)) = \pi\ell_1/p, \quad r(\alpha(y)) = \pi\ell_2/q, \quad r(\alpha(x)\alpha(y)) = \pi\ell_3/r,$$

where ℓ_i is even if $\varepsilon_i = 1$, ℓ_i is odd if $\varepsilon_i = -1$, and $0 < \ell_1 < p$, $0 < \ell_2 < q$, $0 < \ell_3 < r$.

After conjugation, we may assume that $\alpha(x) = e^{\pi i \ell_1/p}$. The quaternions $\alpha(y)$ and $\alpha(x)\alpha(y)$ should lie in their respective conjugacy classes, $S_2 = r^{-1}(\pi\ell_2/q)$ and $S^3 = r^{-1}(\pi\ell_3/r)$. On the other hand, $\alpha(x)\alpha(y)$ lies in $\alpha(x) \cdot S_2$, therefore, in order for $\alpha(x)$ and $\alpha(y)$ to define a representation, the intersection $\alpha(x) \cdot S_2 \cap S_3$ must be non-empty. Since $\alpha(x) \cdot S_2$ is a 2-sphere centered at $\alpha(x)$, the intersection $\alpha(x) \cdot S_2 \cap S_3$ in S^3 (if non-empty) is a circle. This circle parametrizes a whole collection of representations α' coming together with α such that $r(\alpha'(x)) = r(\alpha(x))$, $r(\alpha'(y)) = r(\alpha(y))$, and $r(\alpha'(x)\alpha'(y)) = r(\alpha(x)\alpha(y))$. In fact, all these representations are conjugate to each other by simultaneous conjugation of $\alpha(x)$ and $\alpha(y)$ by the complex circle $S^1 \subset S^3$. This can be seen from the following technical lemma.

Lemma 6. *Let α and β be irreducible representations of the group $\pi_1(\Sigma(p, q, r))$ in $SU(2)$ such that*

- (1) $\alpha(h) = \beta(h)$ and $\alpha(x) = \beta(x) \in \mathbb{C}$,
- (2) $r(\alpha(y)) = r(\beta(y))$, and
- (3) $r(\alpha(x)\alpha(y)) = r(\beta(x)\beta(y))$.

Then the representations α and β are conjugate to each other, that is there exists a unit quaternion c such that

$$\beta(t) = c \cdot \alpha(t) \cdot c^{-1} \quad \text{for all } t \in \pi_1(\Sigma(p, q, r)).$$

Moreover, the quaternion c may be chosen to be a complex number, $c \in \mathbb{C}$.

Corollary 7. (Compare [6]). The representation space $\mathcal{R}(\Sigma(p, q, r))$ is finite.

The involution $\sigma : \Sigma(p, q, r) \rightarrow \Sigma(p, q, r)$ induces the involution on the fundamental group,

$$\begin{aligned} \sigma_* : \pi_1(\Sigma(p, q, r)) &\rightarrow \pi_1(\Sigma(p, q, r)), \\ h &\mapsto h^{-1}, \quad x \mapsto x^{-1}, \quad y \mapsto xy^{-1}x^{-1}, \quad z \mapsto xyz^{-1}y^{-1}x^{-1}, \end{aligned}$$

see [2], Proposition 12.30, which in turn induces an involution on the corresponding representation space ($[\cdot, \cdot]$ stands for conjugacy class),

$$\sigma^* : \mathcal{R}(\Sigma(p, q, r)) \rightarrow \mathcal{R}(\Sigma(p, q, r)), \quad [\alpha] \mapsto [\alpha'], \quad (8)$$

$$\text{where } \alpha'(t) = \alpha(\sigma_*(t)), \quad t \in \pi_1(\Sigma(p, q, r)). \quad (9)$$

Lemma 8. If $\alpha' : \pi_1(\Sigma(p, q, r)) \rightarrow \text{SU}(2)$ is a representation defined by the formula (9) then there exists an element $\rho \in \text{SU}(2)$ such that $\rho^2 = -1$ and

$$\alpha'(t) = \rho \cdot \alpha(t) \cdot \rho^{-1} \quad \text{for all } t \in \pi_1(\Sigma(p, q, r)).$$

The element ρ is defined uniquely up to multiplication by ± 1 , and the elements ρ corresponding to different representations α are conjugate to each other. In particular, the action (8) on the space $\mathcal{R}(\Sigma(p, q, r))$ of the conjugacy classes of irreducible representations of $\pi_1(\Sigma(p, q, r))$ in $\text{SU}(2)$ is trivial.

Proof. After conjugation we may assume that $\alpha(x)$ is a unit complex number. Then $\alpha'(x) = \alpha(\sigma_*(x)) = \alpha(x^{-1}) = \alpha(x)^{-1}$ is a complex number as well. Let us consider another representation, β , defined as a conjugate of α by the unit quaternion j ,

$$\beta(t) = j^{-1} \cdot \alpha'(t) \cdot j \quad \text{for all } t \in \pi_1(\Sigma(p, q, r)).$$

Next we want to verify that the representations α and β satisfy the conditions of Lemma 6. We have

$$\begin{aligned} \beta(x) &= j^{-1} \cdot \alpha'(x) \cdot j \\ &= j^{-1} \cdot \alpha(x)^{-1} \cdot j \\ &= j^{-1} \cdot \overline{\alpha(x)} \cdot j \\ &= j^{-1} \cdot j \cdot \alpha(x), \quad \text{since } \alpha(x) \in \mathbb{C}, \\ &= \alpha(x). \end{aligned}$$

Note that, for any unit quaternion a , we have $r(a) = r(\bar{a}) = r(a^{-1})$. Therefore,

$$\begin{aligned} r(\beta(y)) &= r(j^{-1} \cdot \alpha'(y) \cdot j) = r(\alpha'(y)) \\ &= r(\alpha(xy^{-1}x^{-1})) = r(\alpha(x)\alpha(y)^{-1}\alpha(x)^{-1}) \\ &= r(\alpha(y)^{-1}) = r(\alpha(y)), \end{aligned}$$

and similarly,

$$\begin{aligned} r(\beta(x)\beta(y)) &= r(j^{-1} \cdot \alpha'(x)\alpha'(y) \cdot j) = r(\alpha'(x)\alpha'(y)) \\ &= r(\alpha(x)^{-1}\alpha(x)\alpha(y)^{-1}\alpha(x)^{-1}) = r((\alpha(x)\alpha(y))^{-1}) \\ &= r(\alpha(x)\alpha(y)). \end{aligned}$$

Since $r(\alpha(h))$ is equal to $r(\beta(h))$ we can apply Lemma 6 to the representations α and β to find a complex number $c \in \mathbb{C}$ such that $\beta(t) = c \cdot \alpha(t) \cdot c^{-1}$ for all $t \in \pi_1(\Sigma(p, q, r))$. Since $\beta(t) = j^{-1} \cdot \alpha'(t) \cdot j = c \cdot \alpha(t) \cdot c^{-1}$, we get that $\alpha'(t) = \rho \cdot \alpha(t) \cdot \rho^{-1}$ with $\rho = jc$.

If we apply σ^* twice to the representation α , we will get α again because σ^* is an involution. Therefore, α is conjugate to itself by the element ρ^2 . But the representation α is irreducible, therefore, $\rho^2 = \pm 1$. The following easy computation shows that ρ^2 is in fact -1 ,

$$\rho^2 = jcjc = jj\bar{c}c = -|c|^2 = -1.$$

The uniqueness of ρ up to ± 1 follows from the irreducibility of α . The fact that the elements ρ defined by different representations are conjugate follows from the fact that $\text{tr}(jc) = 0$ for any complex number c . $\square$

4. The definition of λ^ρ . Let $\Sigma = \Sigma(p, q, r)$ be a Brieskorn homology sphere with a σ -invariant Heegaard splitting

$$\Sigma = M_1 \cup_{M_0} M_2 \tag{10}$$

constructed in (6). According to Lemma 8, for any representation $\alpha : \pi_1(\Sigma(p, q, r)) \rightarrow \text{SU}(2)$, there exists an element $\rho \in \text{SU}(2)$ such that $\alpha \circ \sigma_* = \rho \alpha \rho^{-1}$ and $\rho^2 = -1$. Generally speaking, ρ depends on α but it follows from Lemma 8 that the elements ρ corresponding to different α 's are conjugate to each other. Therefore, the following definition makes sense.

For each manifold M_i , $i = 0, 1, 2$, in (10) we define the so called ρ -invariant representation space,

$$\begin{aligned} \mathcal{R}^\rho(M_i) &= \{ \alpha : \pi_1(M_i) \rightarrow \text{SU}(2) \mid \alpha \text{ is irreducible and} \\ &\quad \alpha(\sigma_*(t)) = \rho \alpha(t) \rho^{-1}, \quad t \in \pi_1(M_i) \} / \text{ad } \text{SU}(2)^\rho, \end{aligned} \tag{11}$$

as the space of all irreducible ρ -invariant representations modulo the adjoint action of the 1-dimensional Lie group $\text{SU}(2)^\rho \subset \text{SU}(2)$ consisting of all $g \in \text{SU}(2)$ such that $\rho g = \pm g \rho$. Note that whenever $\rho' = \pm h \rho h^{-1}$, $h \in \text{SU}(2)$, we have that $\mathcal{R}^\rho(M_i) = \mathcal{R}^{\rho'}(M_i)$, hence each of the spaces (11) only depends on the conjugacy class of ρ and not on ρ itself. We do not need to define a ρ -invariant representation space $\mathcal{R}^\rho(\Sigma(p, q, r))$ because all the representations of $\pi_1(\Sigma(p, q, r))$ in $\text{SU}(2)$ are ρ -invariant, see Lemma 8, and therefore $\mathcal{R}^\rho(\Sigma(p, q, r)) = \mathcal{R}(\Sigma(p, q, r))$.

Due to van Kampen's theorem, we have the following commutative diagrams of the fundamental groups

$$\begin{array}{ccc} \pi_1(\Sigma(p, q, r)) & \longleftarrow & \pi_1(M_1) \\ \uparrow & & \uparrow \\ \pi_1(M_2) & \longleftarrow & \pi_1(M_0) \end{array}$$

and of the ρ -invariant representation spaces

$$\begin{array}{ccc} \mathcal{R}(\Sigma(p, q, r)) & \longrightarrow & \mathcal{R}^\rho(M_1) \\ \downarrow & & \downarrow \\ \mathcal{R}^\rho(M_2) & \longrightarrow & \mathcal{R}^\rho(M_0) \end{array}$$

The maps in the latter diagram are injective, so we can think about $\mathcal{R}^\rho(\Sigma(p, q, r))$ as the intersection of $\mathcal{R}^\rho(M_1)$ and $\mathcal{R}^\rho(M_2)$ inside $\mathcal{R}^\rho(M_0)$.

Proposition 9. *The ρ -invariant representation varieties $\mathcal{R}^\rho(M_i)$, $i = 0, 1, 2$, are smooth manifolds of dimensions $\dim \mathcal{R}^\rho(M_i) = 2g - 1$, $i = 1, 2$, and $\dim \mathcal{R}^\rho(M_0) = 4g - 2$ where g is the genus of a σ -invariant Heegaard splitting (10). The maps in the second diagram are embeddings. The submanifolds $\mathcal{R}^\rho(M_1)$ and $\mathcal{R}^\rho(M_2)$ of $\mathcal{R}^\rho(M_0)$ intersect transversally, and their intersection is in one-to-one correspondence with $\mathcal{R}(\Sigma(p, q, r))$.*

The manifolds $\mathcal{R}^\rho(M_i)$, $i = 0, 1, 2$, can be oriented as follows. We start with $\mathcal{R}^\rho(M_0)$. Since the quotient M_0/σ is a 2-sphere and the branching set consists of $2n$ points, $n = g + 1$, one can choose generators $a_1, b_1, \dots, a_g, b_g$ in $\pi_1(M_0)$ such that $\sigma_*(a_i) = a_i^{-1}$, $\sigma_*(b_i) = b_i^{-1}$. Then $\mathcal{R}^\rho(M_0) \subset \text{Hom}^\rho(F_{2g}, \text{SU}(2))/\text{ad } \text{SU}(2)^\rho$ where F_{2g} is a free group on the generators $a_1, b_1, \dots, a_g, b_g$, and

$$\begin{aligned} \text{Hom}^\rho(F_{2g}, \text{SU}(2)) &= \{ \alpha : F_{2g} \rightarrow \text{SU}(2) \mid \\ &\quad \alpha(a_i)^{-1} = \rho \alpha(a_i) \rho^{-1}, \alpha(b_i)^{-1} = \rho \alpha(b_i) \rho^{-1}, i = 1, \dots, g \}. \end{aligned}$$

Lemma 10. *Let $\rho \in \text{SU}(2)$ be such that $\rho^2 = -1$. Then the subset S^ρ of $\text{SU}(2)$ consisting of all $a \in \text{SU}(2)$ such that $a^{-1} = \rho a \rho^{-1}$ is a two-dimensional sphere $S^2 \subset \text{SU}(2)$.*

Proof. Any element ρ of $\text{SU}(2)$ with $\rho^2 = -1$ has zero trace. Therefore, there exists $x \in \text{SU}(2)$ such that $\rho = x j x^{-1}$ (remember that we identify $\text{SU}(2)$ with the group of unit quaternions), and then the map $a \mapsto x^{-1} a x$ establishes a diffeomorphism between S^j and S^ρ . Now, S^j consists of all $b \in \text{SU}(2)$ such that $\bar{b} j = j b$ and $|b|^2 = 1$,

hence $b = u + vi + wk$ for some real u, v, w with $u^2 + v^2 + w^2 = 1$. This of course determines a 2-sphere. $\square$

Our choice of $a_1, b_1, \dots, a_g, b_g$ establishes a diffeomorphism

$$\mathrm{Hom}^\rho(F_{2g}, \mathrm{SU}(2)) = (S^2)^{2g}.$$

By choosing an orientation on S^2 , we orient $\mathrm{Hom}^\rho(F_{2g}, \mathrm{SU}(2))$ as a product. This also orients $\mathcal{R}(M_0)$ by the usual “base-fiber” convention. Note that the orientation on $\mathcal{R}^\rho(M_0)$ is independent of our choices of orientations on S^2 and $\mathrm{SU}(2)^\rho$ or our choice of a basis $a_1, b_1, \dots, a_g, b_g$.

The orientation on $\mathcal{R}^\rho(M_1)$ can be defined as follows. First, we choose a basis $x_1, \dots, x_g$ of the free group $\pi_1(M_1)$ so that $\sigma_*(x_i) = x_i^{-1}$, $i = 1, \dots, g$. Then $\mathcal{R}^\rho(M_1) \subset \mathrm{Hom}^\rho(F_g, \mathrm{SU}(2))/\mathrm{ad}\mathrm{SU}(2)^\rho$, where F_g is a free group on the generators $x_1, \dots, x_g$, and

$$\begin{aligned} \mathrm{Hom}^\rho(F_g, \mathrm{SU}(2)) &= \{ \alpha : F_g \rightarrow \mathrm{SU}(2) \mid \alpha(\sigma_*(x)) = \rho\alpha(x)\rho^{-1} \} \\ &= \{ \alpha : F_g \rightarrow \mathrm{SU}(2) \mid \alpha(x_i)^{-1} = \rho\alpha(x_i)\rho^{-1}, \ i = 1, \dots, g \}. \end{aligned}$$

Our choice of $x_1, \dots, x_n$ establishes a diffeomorphism

$$\mathrm{Hom}^\rho(F_g, \mathrm{SU}(2)) = (S^2)^g.$$

We choose an orientation on S^2 and orient $\mathrm{Hom}^\rho(F_g, \mathrm{SU}(2))$ as a product. Note that this orientation changes by $(-1)^g$ when the orientation on S^2 is changed; however, it does not depend on the choice of a basis in $\pi_1(M_1) = F_g$. The orientation of $\mathrm{Hom}^\rho(F_g, \mathrm{SU}(2))$ and an orientation of $\mathrm{SU}(2)^\rho$ orient $\mathcal{R}(M_1)$.

The ρ -invariant representation space $\mathcal{R}^\rho(M_2)$ is oriented in a completely similar way, the orientation of $\mathrm{SU}(2)^\rho$ having already been fixed.

If the orientations on S^2 or $\mathrm{SU}(2)^\rho$ are changed, the spaces $\mathcal{R}(M_1)$ and $\mathcal{R}(M_2)$ change their orientations simultaneously; therefore, their algebraic intersection number in $\mathcal{R}^\rho(M_0)$ does not change (though *a priori* it may depend on the chosen Heegaard decomposition). We define our λ^ρ -invariant as one half of this algebraic intersection number, so that

$$\lambda^\rho(\Sigma(p, q, r)) = 1/2 \cdot \sum_{\alpha \in \mathcal{R}(\Sigma)} \varepsilon_\alpha \quad (12)$$

where ε_α equals ± 1 depending on whether the orientations on the tangent spaces $T_\alpha \mathcal{R}^\rho(M_1) \oplus T_\alpha \mathcal{R}^\rho(M_2)$ and $T_\alpha \mathcal{R}^\rho(M_0)$ agree.

Proposition 11. *For any Brieskorn homology sphere $\Sigma(p, q, r)$ the invariant $\lambda^\rho(\Sigma(p, q, r))$ is well-defined, in particular, it does not depend on the choice of a σ -invariant Heegaard splitting. Moreover,*

$$\lambda^\rho(\Sigma(p, q, r)) = 1/8 \cdot \mathrm{sign} \, k(p, q, r)$$

where sign stands for the knot signature.

This proposition is proven below by pushing invariant representations down to a knot $k(p, q, r)$ complement in S^3 and applying a result from [8] on trace-free $SU(2)$ -representations of knot groups.

Thus our definition of λ^ρ -invariant is modelled on the Casson's definition of the λ -invariant; the difference is that we use ρ -invariant representation spaces. Surprisingly enough, we get an invariant which is *different* from λ . For example, $\lambda(\Sigma(2, 3, 5)) = \lambda(\Sigma(2, 3, 7)) = 1$ while $\lambda^\rho(\Sigma(2, 3, 5)) = -1$ and $\lambda^\rho(\Sigma(2, 3, 7)) = 1$.

5. Casson-Lin invariant. Let B_n be the braid group of rank n with the standard generators $\beta_1, \dots, \beta_{n-1}$ represented in a free group F_n on symbols $x_1, \dots, x_n$ as follows:

$$\begin{aligned} \beta_i : \quad x_i &\mapsto x_i x_{i+1} x_i^{-1} \\ x_{i+1} &\mapsto x_i \\ x_j &\mapsto x_j, \quad \text{if } j \neq i, i+1. \end{aligned}$$

If $\beta \in B_n$ then the automorphism of F_n representing β maps each x_i to a conjugate of some x_j and preserves the product $x_1 \cdots x_n$.

Let $k \subset S^3$ be a knot represented as the closure of a braid $\beta \in B_n$. Let us fix an embedding of k into S^3 as shown in Figure 2, the sphere S separating S^3 in two 3-balls, B_1 and B_2 , with the braid β inside B_1 and n untangled arcs inside B_2 . The fundamental group $\pi_1(K)$ of the knot k complement $K = S^3 - nbd(k)$, has the presentation

$$\pi_1(K) = \langle x_1, \dots, x_n \mid x_i = \beta(x_i), \ i = 1, \dots, n \rangle,$$

the generators $x_1, \dots, x_n$ being represented by the meridians of β .

The knot complement K can be now decomposed as $K = M'_1 \cup_{M'_0} M'_2$ where $M'_0 = S \cap K$ and $M'_i = B_i \cap K$, $i = 1, 2$. The manifolds M'_1 and M'_2 are handlebodies of genus n , and M'_0 is a 2-sphere with $2n$ small 2-discs removed around the points $P_1, \dots, P_{2n}$, see Figure 2. Due to van Kampen's theorem, we get the following commutative diagram of fundamental groups

$$\begin{array}{ccc} \pi_1(K) & \longleftarrow & \pi_1(M'_1) \\ \uparrow & & \uparrow \\ \pi_1(M'_2) & \longleftarrow & \pi_1(M'_0) \end{array}$$

where

$$\begin{aligned} \pi_1(M'_0) &= \langle x_1, \dots, x_n, y_1, \dots, y_n \mid x_1 \cdots x_n = y_1 \cdots y_n \rangle, \\ \pi_1(M'_1) &= \langle x_1, \dots, x_n \mid \rangle, \quad \pi_1(M'_2) = \langle y_1, \dots, y_n \mid \rangle \end{aligned}$$

are isomorphic to free groups. To continue an analogy with the definition of the Casson invariant, we need to define $\mathrm{SU}(2)$ -representation spaces of the groups $\pi_1(K)$ and $\pi_1(M_i)$, $i = 0, 1, 2$, and compute the corresponding intersection number. To make things work, we impose the extra condition on the representations that all the meridians go to trace-free matrices in $\mathrm{SU}(2)$. Thus, following [8], we define

$$\begin{aligned}\hat{H}_n &= \{ \alpha : \pi_1(M'_0) \rightarrow \mathrm{SU}(2) \mid \alpha \text{ is irreducible, } \mathrm{tr} x_i = \mathrm{tr} y_i = 0 \} / \mathrm{ad} \mathrm{SU}(2), \\ \hat{\Gamma}_\beta &= \{ \alpha : \pi_1(M'_1) \rightarrow \mathrm{SU}(2) \mid \alpha \text{ is irreducible, } \mathrm{tr} x_i = 0 \} / \mathrm{ad} \mathrm{SU}(2), \\ \hat{\Lambda}_n &= \{ \alpha : \pi_1(M'_2) \rightarrow \mathrm{SU}(2) \mid \alpha \text{ is irreducible, } \mathrm{tr} y_i = 0 \} / \mathrm{ad} \mathrm{SU}(2), \\ \mathcal{R}^0(K) &= \{ \alpha : \pi_1(K) \rightarrow \mathrm{SU}(2) \mid \alpha \text{ is irreducible, } \mathrm{tr} x_i = 0 \} / \mathrm{ad} \mathrm{SU}(2).\end{aligned}$$

The first three are smooth manifolds of dimensions $\dim \hat{H}_n = 4n - 6$, and $\dim \hat{\Gamma}_\beta = \dim \hat{\Lambda}_n = 2n - 3$. The commutative diagram of fundamental groups induces the following commutative diagram

$$\begin{array}{ccc}\mathcal{R}^0(K) & \longrightarrow & \hat{\Gamma}_\beta \\ \downarrow & & \downarrow \\ \hat{\Lambda}_n & \longrightarrow & \hat{H}_n\end{array}$$

In particular, the irreducible trace-free representations of $\pi_1(K)$ in $\mathrm{SU}(2)$ are in one-to-one correspondence with the intersection points of $\hat{\Lambda}_n$ with $\hat{\Gamma}_\beta$ in $\hat{H}_n$. The manifolds $\hat{\Lambda}_n$, $\hat{\Gamma}_\beta$, and $\hat{H}_n$ are naturally oriented, see [8], and (possibly after a perturbation to make the intersection transversal) one can define

$$h(k) = \sum_{\alpha \in \mathcal{R}^0(K)} \varepsilon'_\alpha \quad (13)$$

where $\varepsilon'_\alpha = \pm 1$ is a sign obtained by comparing the orientations on $T_\alpha \hat{\Lambda}_n \oplus T_\alpha \hat{\Gamma}_\beta$ and $T_\alpha \hat{H}_n$. The invariant $h(k)$ only depends on the knot k and not on the choices in the definition. X.-S. Lin in [8] further proves that $h(k) = 1/2 \cdot \mathrm{sign} k$.

6. Representations of a knot $k(p, q, r)$ complement. Throughout this subsection $\Sigma = \Sigma(p, q, r)$ is a Brieskorn homology sphere, and $k = k(p, q, r)$ is the corresponding Montesinos knot.

Let $E \rightarrow \Sigma$ be a (necessarily trivial) $\mathrm{SU}(2)$ -vector bundle over Σ . Let us fix a Riemannian metric on Σ and consider the Banach manifold

$$\mathcal{B}^* = \mathcal{A}^* / \mathcal{G}$$

of the L^2_2 -gauge equivalence classes of irreducible L^2_1 -connections in E . It is a classical result in differential geometry that the holonomy map establishes a one-to-one

correspondence between the gauge equivalence classes of irreducible flat connections and the conjugacy classes of irreducible $\mathrm{SU}(2)$ -representations of $\pi_1(\Sigma)$.

The involution σ can be lifted to a bundle endomorphism of E . Any endomorphism of E clearly induces an action on $\mathcal{A}^*$ by pull-back, and an action on $\mathcal{B}^*$ as well. Since any two liftings of σ differ by a gauge transformation, we have a well-defined action $\sigma^* : \mathcal{B}^* \rightarrow \mathcal{B}^*$. Denote by $\mathcal{B}^\sigma$ the space of connections invariant with respect to σ^* .

Let $\rho \in \mathrm{SU}(2)$ be such that $\rho^2 = -1$. The formula

$$(x, \xi) \mapsto (\sigma(x), \rho\xi\rho^{-1}) \quad (14)$$

defines a lifting of $\sigma : \Sigma \rightarrow \Sigma$ on E , which will again be denoted by $\rho : E \rightarrow E$. Let $\mathcal{B}^\rho \subset \mathcal{B}^\sigma$ consist of the gauge equivalence classes of irreducible connections A in E such that $\rho^*A = A$. Due to Lemma 8, all irreducible flat connections on Σ belong to $\mathcal{B}^\rho$; in particular, $\mathcal{B}^\rho$ is non-empty. The following three lemmas are easy corollaries of Propositions 1 and 17 and Theorem 18 in [20].

Lemma 12. *For any $\rho, \rho' \in \mathrm{SU}(2)$ such that $\rho^2 = \rho'^2 = -1$, $\mathcal{B}^\rho = \mathcal{B}^{\rho'}$. Furthermore, as a set, $\mathcal{B}^\rho$ is bijective to $\mathcal{A}^\rho/\overline{\mathcal{G}}^\rho$ where $\overline{\mathcal{G}} = \mathcal{G}/\pm 1$ and $\mathcal{A}^\rho = \{A \in \mathcal{A}^* \mid \rho^*A = A\}$.*

The quotient of $\Sigma(p, q, r)$ by σ is S^3 . Since $\rho \neq \pm 1$, it is impossible to define the quotient bundle of E over S^3 . One can though define it away from the image of the fixed point set of σ , that is, on the knot complement $K = S^3 \setminus k$. Given an irreducible flat ρ -invariant $\mathrm{SU}(2)$ -connection $A \in \mathcal{A}^\rho$, its push-down A' is an irreducible flat $\mathrm{SO}(3)$ -connection on K . In other words, A' is an irreducible flat $\mathrm{SO}(3)$ -connection singular along k in the sense of P. Kronheimer and T. Mrowka, see [7].

Lemma 13. *The $\mathrm{SO}(3)$ -connection A' has holonomy $1/2$ around k .*

According to $H^1(K; \mathbb{Z}/2) = \mathbb{Z}/2$ and $H^2(K; \mathbb{Z}/2) = 0$, there are two different ways to lift the $\mathrm{SO}(3)$ -connection A' to a (singular) $\mathrm{SU}(2)$ -connection. Both $\mathrm{SU}(2)$ -liftings have holonomy $1/4$ around k ; in other words, their holonomy around k is trace-free.

Lemma 14. *Through the push-down map, the representation space $\mathcal{R}(\Sigma)$ is in one-to-two correspondence with the representation space $\mathcal{R}^0(K)$ of irreducible trace-free $\mathrm{SU}(2)$ -representations of $\pi_1(K)$.*

Similar one-to-two identifications through push-down hold for the pairs $\mathcal{R}^\rho(M_0)$ and $\hat{H}_n$, $\mathcal{R}^\rho(M_1)$ and $\hat{\Gamma}_\beta$, $\mathcal{R}^\rho(M_2)$ and $\hat{\Lambda}_n$, where $g = n - 1$ is the genus of the σ -invariant Heegaard splitting, see Figure 2.

Proof of Proposition 11. Remember that throughout this subsection, $\Sigma = \Sigma(p, q, r)$ and $k = k(p, q, r)$. The invariants $\lambda^\rho(\Sigma)$ and $h(k) = 1/2 \cdot \mathrm{sign} k$ were defined in (12) and (13) as an algebraic count of points in the corresponding (finite) representation

spaces, $\mathcal{R}(\Sigma)$ and $\mathcal{R}^0(K)$. These spaces are in one-to-two correspondence by Lemma 14. A routine comparison of the orientations shows that $\varepsilon_\alpha = \varepsilon'_{\alpha'}$ where α' is a push-down of α . Therefore,

$$\begin{aligned}\lambda^\rho(\Sigma) &= 1/2 \cdot \sum \varepsilon_\alpha \\ &= 1/4 \cdot \sum \varepsilon'_{\alpha'} \\ &= 1/8 \cdot \text{sign } k.\end{aligned}$$

This completes the proof of Proposition 11. $\square$

3. GAUGE THEORY FOR BRIESKORN HOMOLOGY SPHERES

Our next step is to give an “analytic” definition of the invariant λ^ρ . In [19], C. Taubes gave a gauge-theoretical interpretation of the Casson’s λ -invariant in terms of (what later became known as) Floer homology. Our gauge-theoretical interpretation of λ^ρ will follow the same lines with the only difference that everything will be “ ρ -invariant”.

1. Casson invariant via gauge theory. Let us recall shortly the Taubes construction [19] for a homology 3-sphere Σ . Let $E \rightarrow \Sigma$ be a trivial $\text{SU}(2)$ -bundle, and $\mathcal{B}^* = \mathcal{A}^*/\mathcal{G}$ the space of gauge equivalence classes of irreducible connections in E . It is a classical differential geometry result that flat connections in E (modulo gauge equivalence) are in one-to-one correspondence with $\text{SU}(2)$ -representations of $\pi_1(\Sigma)$ (modulo conjugation). Given two irreducible representations α, β , we may think of them as points in $\mathcal{A}^*$ or $\mathcal{B}^*$.

For any $A \in \mathcal{A}^*$, define the following elliptic differential operator

$$K_A = \begin{vmatrix} 0 & d_A^* \\ d_A & *d_A \end{vmatrix} : (\Omega^0 \oplus \Omega^1)(\Sigma, \mathfrak{su}(2)) \rightarrow (\Omega^0 \oplus \Omega^1)(\Sigma, \mathfrak{su}(2)), \quad (15)$$

where d_A stands for the operator of covariant differentiation with respect to A , and $*$ is the Hodge operator associated with a Riemannian metric on Σ . In a proper Sobolev completion of $(\Omega^0 \oplus \Omega^1)(\Sigma, \mathfrak{su}(2))$, the operator K_A is Fredholm. Let $A(t)$, $0 \leq t \leq 1$, be a path in $\mathcal{A}^*$ connecting α with β , so that $A(0) = \alpha$, $A(1) = \beta$. Associated with $A(t)$ is a path of Fredholm operators, $K_{A(t)}$. Let us assume that $\ker K_{A(0)} = \ker K_{A(1)}$, as is the case for a Brieskorn homology sphere $\Sigma = \Sigma(p, q, r)$. We define $sf(\alpha, \beta)$ as the spectral flow of the path $K_{A(t)}$ between α and β , see [1]. This number is well-defined modulo 8, and Taubes defines an infinite dimensional generalization $\chi(\Sigma)$ of the Euler characteristic as

$$\chi(\Sigma) = \varepsilon_\alpha \cdot \sum_{\beta \in \mathcal{R}(\Sigma)} (-1)^{sf(\alpha, \beta)} \quad (16)$$

where ε_α is figured out from the spectral flow between the trivial connection θ and α (the number $\chi(\Sigma)$ turns out to be independent of α). It is proven in [19] that $\lambda(\Sigma) = 1/2 \cdot \chi(\Sigma)$. Later, A. Floer showed in [5] that there are well-defined $\mathbb{Z}/8$ -graded instanton homology groups $I_*(\Sigma)$ such that $\chi(\Sigma) = \chi(I_*(\Sigma))$.

Let now $\Sigma = \Sigma(p, q, r)$ be a Brieskorn homology sphere. R. Fintushel and R. Stern showed in [6] (see also Proposition 15 below) that for any pair $\alpha, \beta \in \mathcal{R}(\Sigma(p, q, r))$,

$$sf(\alpha, \beta) = 0 \pmod{2},$$

and that all the signs involved in computing $\chi(\Sigma(p, q, r))$ are positive, so χ simply counts the irreducible representations.

2. Floer index for Brieskorn homology spheres. Let α and β be irreducible flat connections in a trivial $SU(2)$ -bundle E over $\Sigma = \Sigma(p, q, r)$. By pull-back, we can extend E to a trivial bundle (which we also denote by E) over the infinite cylinder $\Sigma \times \mathbb{R}$. Let us choose a σ -invariant Riemannian metric on Σ , and the corresponding product metric on $\Sigma \times \mathbb{R}$. Let further $\rho : E \rightarrow E$ be the lifting of σ defined in (14), and $A(t) \in \mathcal{A}^\rho$ a path of connections forming an invariant connection A over $\Sigma \times \mathbb{R}$ vanishing in the $\mathbb{R}$ -direction and equal to respectively α and β near the ends of $\Sigma \times \mathbb{R}$. Denote by d_A the operator of covariant differentiation with respect to A .

The (relative) Floer index of the pair (α, β) equals the Fredholm index of the following elliptic complex, see [1],

$$\Omega^0(\Sigma \times \mathbb{R}, \text{ad } E) \xrightarrow{d_A} \Omega^1(\Sigma \times \mathbb{R}, \text{ad } E) \xrightarrow{d_A^-} \Omega_+^2(\Sigma \times \mathbb{R}, \text{ad } E) \quad (17)$$

where $\text{ad } E$ is the adjoint bundle of E over $\Sigma \times \mathbb{R}$, and $d_A^- = P_- \circ d_A$ where P_- is the projection onto the self dual forms with respect to the fixed product metric on $\Sigma \times \mathbb{R}$. The Sobolev norms on the spaces in (17) are specified as usual, see [5].

The product metric and the connection A on $\Sigma \times \mathbb{R}$ are ρ -invariant, therefore, one can define the following ρ -invariant elliptic subcomplex of (17),

$$\Omega^0(\Sigma \times \mathbb{R}, \text{ad } E)^\rho \xrightarrow{d_A} \Omega^1(\Sigma \times \mathbb{R}, \text{ad } E)^\rho \xrightarrow{d_A^-} \Omega_+^2(\Sigma \times \mathbb{R}, \text{ad } E)^\rho \quad (18)$$

Proposition 15. *For any irreducible flat connections α, β on $\Sigma = \Sigma(p, q, r)$, the index of the elliptic complex (17) is even. In fact, it equals twice the index of the ρ -invariant complex (18).*

In order to prove the proposition we need the following two technical lemmas.

Lemma 16. *There exist a Riemannian metric on $\Sigma(p, q, r)$ and an almost complex structure J on $\Sigma(p, q, r) \times \mathbb{R}$ such that*

- (1) *The product metric on $\Sigma \times \mathbb{R}$ is $(\sigma \times 1)$ -invariant;*
- (2) *J is compatible with this product metric; and*
- (3) *The involution $\sigma \times 1$ is anti-holomorphic with respect to J , that is $(\sigma \times 1)_* J = -J(\sigma \times 1)_*$ on the tangent bundle.*

Proof. Let us consider the algebraic variety

$$V(p, q, r) = \{ (x, y, z) \in \mathbb{C}^3 \mid x^p + y^q + z^r = 0 \} \subset \mathbb{C}^3,$$

which is a non-singular complex surface except perhaps at the origin. The homology sphere $\Sigma(p, q, r)$ in question is the intersection of $V(p, q, r)$ with the unit 5-sphere S_1^5 ,

$$\Sigma(p, q, r) = V(p, q, r) \cap S_1^5,$$

and the variety $V(p, q, r)$ is in fact a cone over $\Sigma(p, q, r)$ with the vertex at the origin.

Let $V^0(p, q, r)$ be the variety $V(p, q, r)$ with the origin removed. The Riemann metric induced on $V^0(p, q, r)$ from the standard flat metric on $\mathbb{C}^3$ is a cone metric given by

$$ds^2 = dr^2 + r^2 d\theta^2 \quad (19)$$

in the spherical coordinates (r, θ) . Here $d\theta^2 = \sum h_{ij} d\theta_i d\theta_j$ is a metric on $\Sigma(p, q, r)$, and r is the distance from the origin (in $\mathbb{C}^3$). Obviously, both the complex conjugation σ' and the almost complex structure J induced on $V^0(p, q, r)$ from $\mathbb{C}^3$ are compatible with the metric (19).

Let us now form the conformally equivalent metric

$$ds^2/r^2 = dr^2/r^2 + d\theta^2 \quad (20)$$

on $V^0(p, q, r)$. The substitution $r = e^{-\tau}$ gives coordinates in which ds^2/r^2 is the standard product metric

$$d\tilde{s}^2 = d\tau^2 + d\theta^2 \quad (21)$$

on the cylinder $\Sigma(p, q, r) \times \mathbb{R}$. Since the metrics (19) and (21) are conformally equivalent, both σ' and J are compatible with the product metric (21). Moreover, σ' preserves all the spheres S_r^5 , $0 < r < \infty$, in $\mathbb{C}^3$, therefore, it has the form $\sigma' = \sigma \times 1$. Finally, the involution $\sigma \times 1$ is antiholomorphic with respect to J because this is the case in $\mathbb{C}^3$. $\square$

Lemma 17. *There exists a differential operator $\tilde{d}_A^- : \Omega^1(\Sigma \times \mathbb{R}, \text{ad } E) \rightarrow \Omega_+^2(\Sigma \times \mathbb{R}, \text{ad } E)$ such that*

- (1) *The difference $\tilde{d}_A^- - d_A^-$ is a compact operator,*
- (2) *$\text{coker } \tilde{d}_A^- = 0$, and*
- (3) *The operator $\tilde{d}_A^-$ is ρ -invariant.*

Proof. This is essentially the transversality result from [5], Proposition 2c.2. The key difference is that we have to bound ourselves to invariant perturbations which, generally speaking, may be not generic among the perturbations used by Floer. Following [5], Section 1b, consider a collection $S(m)$ of m circles smoothly embedded in

$\mathbb{R}^3$ which intersect precisely at the origin and have the same tangent there. Let

$$\gamma : \bigvee_{i=1}^m S_i^1 \times D^2 \rightarrow \Sigma,$$

be a map which restricts to smooth embeddings $\gamma_\theta : S(m) \rightarrow \Sigma$ for each $\theta \in D^2$ and $\gamma_i : S_i^1 \times D^2 \rightarrow \Sigma$ for each i . By adding extra circles, one can choose γ so that its image in Σ will be σ -invariant. Now, γ defines a family of holonomy maps

$$\begin{aligned} \gamma_\theta : \mathcal{B} &\rightarrow L_m = \mathrm{SU}(2)^m / \mathrm{ad} \mathrm{SU}(2), \\ A &\mapsto (\mathrm{hol}_A(\gamma_1), \dots, \mathrm{hol}_A(\gamma_m)) \end{aligned} \quad (22)$$

Let $C_{sym}^\infty(L_m, \mathbb{R})$ be defined as the set of smooth $\mathrm{ad} \mathrm{SU}(2)$ -invariant real valued functions on $\mathrm{SU}(2)^m$ with the additional property that for any $h \in C_{sym}^\infty(L_m, \mathbb{R})$,

$$h(x_{\tau(1)}, \dots, x_{\tau(m)}) = h(x_1, \dots, x_m)$$

for any permutation τ on m symbols. For example, one can start with an arbitrary $\mathrm{ad} \mathrm{SU}(2)$ -invariant function on $\mathrm{SU}(2)^m$ and take h to be its symmetrization.

Given such γ and h , we define the function

$$h_\gamma : \mathcal{B} \rightarrow \mathbb{R}, \quad h_\gamma(A) = \int_{D^2} h(\gamma_\theta(A)) d^2\theta \quad (23)$$

where $d^2\theta$ is a smooth compactly supported volume form on the interior of D^2 . The following argument shows that this function is invariant with respect to the induced σ^* -action on $\mathcal{B}$:

$$\begin{aligned} h_\gamma(\sigma^* A) &= \int_{D^2} h(\mathrm{hol}_{\sigma^* A}(\gamma_1), \dots, \mathrm{hol}_{\sigma^* A}(\gamma_m)) d^2\theta \\ &= \int_{D^2} h(\mathrm{hol}_A(\sigma(\gamma_1)), \dots, \mathrm{hol}_A(\sigma(\gamma_m))) d^2\theta \\ &= \int_{D^2} h(\mathrm{hol}_A(\gamma_{\tau(1)}), \dots, \mathrm{hol}_A(\gamma_{\tau(m)})) d^2\theta \\ &= \int_{D^2} h(\mathrm{hol}_A(\gamma_1), \dots, \mathrm{hol}_A(\gamma_m)) d^2\theta = h_\gamma(A). \end{aligned}$$

The class of maps we defined in (23) is large enough for the set

$$\{ \mathrm{grad} h_\gamma(A) \mid h_\gamma \text{ defined by (23)} \} \subset T_A \mathcal{B}^*$$

to be dense in the tangent space $T_A \mathcal{B}^*$. Now the proof of Proposition 2c.2 in [5] goes through, which provides a function h_γ such that a linearization of its gradient is a compact invariant perturbation of the operator d_A^- with the desired properties. $\square$

Sketch of the proof of Proposition 15. In Lemma 16 we defined a σ -invariant Riemannian metric on $\Sigma(p, q, r)$ and an almost complex structure J on the manifold $\Sigma(p, q, r) \times \mathbb{R}$ compatible with the product metric such that the involution $\sigma \times 1 : \Sigma \times \mathbb{R} \rightarrow \Sigma \times \mathbb{R}$ is anti-holomorphic with respect to J . We can use J to identify the ± 1 -eigenspaces of the involution induced by $\sigma \times 1$ on the space of 1-forms in (17), and make sure that (possibly after perturbation provided by Lemma 17) $\ker d_A = 0$ and $\operatorname{coker} d_A^- = 0$. This splits the index of the complex (17) and hence the corresponding spectral flow in halves. $\square$

Proposition 18. (Compare [6]). *Let $\Sigma(p, q, r)$ be a Brieskorn homology sphere. Then the Floer homology groups $I_n(\Sigma(p, q, r))$ vanish for n odd.*

Proof. We only need to fix the sign ε_α in (16). It can be achieved by the requirement that for $\varepsilon_\alpha = (-1)^{\eta(\alpha)}$,

$$-3 - \eta(\alpha) = \operatorname{index} D_A^+ - 3(1 + b_2^+)(W),$$

where W is any smooth simply connected non-compact 4-manifold with a single end of the form $\Sigma(p, q, r) \times \mathbb{R}_+$, A is any $\mathrm{SU}(2)$ -connection on W with limiting value α , and

$$D_A^+ = d_A \oplus d_A^+ : \Omega^1(W, \mathfrak{su}(2)) \rightarrow \Omega^0(W, \mathfrak{su}(2)) \oplus \Omega_+^2(W, \mathfrak{su}(2))$$

is the ASD-operator, compare [4]. Let W be the Milnor fiber of the singularity of $f^{-1}(0)$ where $f(x, y, z) = x^p + y^q + z^r$ so that $\partial W = \Sigma(p, q, r)$, see [9]. The involution (5) extends to an antiholomorphic involution on W , and an argument similar to that in the proof of Proposition 15 proves that $\operatorname{index} D_A^+$ is even. On the other hand, one can easily show that modulo 2, $b_2^+ = 1/2 \cdot b_2(W) = 1/2 \cdot (p-1)(q-1)(r-1) = 0$. Therefore, $\varepsilon_\alpha = 1$, and the Floer homology groups of $\Sigma(p, q, r)$ vanish in odd dimensions. $\square$

3. The definition of χ^ρ . Let $\rho : E \rightarrow E$ be the lifting of $\sigma : \Sigma(p, q, r) \rightarrow \Sigma(p, q, r)$ defined by (14). It induces an involution ρ^* on the $\mathfrak{su}(2)$ -valued differential forms on $\Sigma(p, q, r)$. Therefore, for any $A \in \mathcal{A}^\rho$, the differential operator K_A , see (15), can be restricted to the $+1$ -eigenspaces of the involution ρ^* to give a new elliptic differential operator K_A^ρ ,

$$K_A^\rho = \begin{vmatrix} 0 & d_A^* \\ d_A & *d_A \end{vmatrix} : (\Omega^0 \oplus \Omega^1)^\rho(\Sigma, \mathfrak{su}(2)) \rightarrow (\Omega^0 \oplus \Omega^1)^\rho(\Sigma, \mathfrak{su}(2)).$$

For any pair $\alpha, \beta \in \mathcal{R}(\Sigma(p, q, r))$, let us denote by $sf^\rho(\alpha, \beta)$ the spectral flow of a family of operators $K_{A(t)}^\rho$, where $A(t)$ is a path in $\mathcal{A}^\rho$ connecting α to β . The number $sf^\rho(\alpha, \beta)$ is well-defined modulo 4. The invariant $\chi^\rho(\Sigma(p, q, r))$ is now defined by the

formula

$$\chi^\rho(\Sigma) = \varepsilon_\alpha \cdot \sum_{\beta \in \mathcal{R}(\Sigma)} (-1)^{sf^\rho(\alpha, \beta)}, \quad (24)$$

compare to (16). The following result is an easy consequence of Proposition 15.

Proposition 19. *Let $\Sigma = \Sigma(p, q, r)$ be a Brieskorn homology sphere. Then $1/2 \cdot \chi^\rho(\Sigma) = \nu(\Sigma)$ where the invariants $\chi^\rho(\Sigma)$ and $\nu(\Sigma)$ are defined by (24) and (1).*

4. THE INVARIANT λ^ρ VIA GAUGE THEORY

Proposition 20. *Let $\Sigma = \Sigma(p, q, r)$ be a Brieskorn homology sphere, and λ^ρ and χ^ρ the invariant defined by (12) and (24). Then $\lambda^\rho(\Sigma) = 1/2 \cdot \chi^\rho(\Sigma)$.*

This result is a straightforward application of the Taubes argument in [19] in our ρ -invariant setting.

REFERENCES

- [1] M. Atiyah, V. Patodi, I. Singer, *Spectral asymmetry and Riemannian geometry. III*, Math. Proc. Cambridge Phil. Soc. **79** (1976), 71–99.
- [2] G. Burde, H. Zieschang, *Knots*, Walter de Gruyter, 1985.
- [3] A. Casson, J. Harer, *Some homology lens spaces which bound rational homology balls*, Pacific J. Math. **96** (1981), 23–36.
- [4] S. Donaldson, M. Furuta, D. Kotschick, *Floer homology groups in Yang-Mills theory*, Preprint.
- [5] A. Floer, *An instanton-invariant for 3-manifolds*, Comm. Math. Phys. **118** (1988), 215–240.
- [6] R. Fintushel, R. Stern, *Instanton homology groups of Seifert fibered homology three spheres*, Proc. London Math. Soc. **61** (1990), 109–137.
- [7] P. Kronheimer, T. Mrowka, *Gauge theory for embedded surfaces. I*, Topology **32** (1993), 773–826.
- [8] X.-S. Lin, *A knot invariant via representation spaces*, J. Diff. Geom. **35** (1992), 337–357.
- [9] J. Milnor, *Singular points of complex hypersurfaces*, Annals of Math. Studies **91**, Princeton, 1968.
- [10] J. Montesinos, *Variedades de Seifert que son recubridores ciclicos ramificados de dos hojas*, Bol. Soc. Mat. Mexicana **18** (1973), 1–32.
- [11] D. Mullins, *The generalized Casson invariant for 2-fold branched covers of S^3 and the Jones polynomial*, Topology **32** (1993), 419–438.
- [12] W. Neumann, J. Wahl, *Casson invariant of link of singularities*, Comm. Math. Helv. **65** (1990), 58–78.
- [13] W. Neumann, *An invariant of plumbed homology spheres*, Lecture Notes in Math. **788**, Springer-Verlag, 1980, 125–144.
- [14] W. Neumann, F. Raymond, *Seifert manifolds, plumbing, μ -invariant and orientation reversing maps*, Lecture Notes in Math. **664**, Springer-Verlag, 1978, 163–196.
- [15] N. Saveliev, *Floer homology and 3-manifold invariants*, Ph.D. Thesis, University of Oklahoma, Norman OK, 1995.
- [16] N. Saveliev, *Dehn surgery along torus knots*, Topology and Its Appl. (to appear), 1996.
- [17] L. Siebenmann, *On vanishing of the Rohlin invariant and nonfinitely amphicheiral homology 3-spheres*, Lecture Notes in Math. **788**, Springer-Verlag, 1980, 172–222.
- [18] R. Stern, *Some more Brieskorn spheres which bound contractible manifolds*, Notices Amer. Math. Soc. **25** (1978), A448.
- [19] C. Taubes, *Casson’s invariant and gauge theory*, J. Diff. Geom. **31** (1990), 547–599.
- [20] S. Wang, *Moduli spaces over manifolds with involutions*, Math. Ann. **296** (1993), 119–138.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109

E-mail address: saveliev@math.lsa.umich.edu